

Numerical Solution of Ordinary Differential Equations

Chapter 25

Differential Equations




- Differential equations play a fundamental role in engineering. Many physical phenomena are best formulated in terms of their rate of change:

$$\frac{dv}{dt} = g - \frac{c}{m}v$$

v - dependent variable
 t - independent variable

- Equations which are composed of an *unknown function* and its *derivatives* are called ***differential equations***.
- **One** independent variable → ***ordinary differential equation*** (or ***ODE***)
- **Two or more** independent variables → ***partial diff. equation*** (or ***PDE***)
- ***A first order equation*** includes a first derivative as its highest derivative
- ***Second order equation*** includes second derivative
- Higher order equations can be reduced to a system of first order equations, by redefining the variables.

ODEs and Engineering Practice



Swinging pendulum

$$\frac{d^2\theta}{dt^2} + \frac{g}{l} \sin \theta = 0$$

A second-order nonlinear ODE.

Falling parachutist problem

Solving Ordinary Differential Equations (ODEs)

- This chapter is devoted to solving ordinary differential equations (ODEs) of the form

$$\frac{dy}{dx} = f(x, y)$$

New value = old value + slope * (step_size)

$$y_{i+1} = y_i + \phi * h$$



Euler's Method

- First derivative provides a direct estimate of the **slope** at x_i :

$$\phi = f(x_i, y_i) \quad (\text{diff.equ.evaluated at } x_i \text{ and } y_i)$$

then,

$$y_{i+1} = y_i + f(x_i, y_i)h$$



Error Analysis for Euler's Method

- Numerical solutions of ODEs involves two types of error:
 - Truncation* error
 - Local** truncation error
 - Propagated truncation error
 The sum of the two is the *total or global* truncation error
 - Round-off* errors (due to limited digits in representing numbers in a computer)
- We can use Taylor series to quantify the **local truncation error** in Euler's method.

Given $y' = f(x, y)$

$$y_{i+1} = y_i + y_i' h + \frac{y_i''}{2!} h^2 + \dots + \frac{y_i^{(n)}}{n!} h^n + R_n$$

$$E_a = \frac{f'(x_i, y_i)}{2!} h^2 + R_3$$

$$y_{i+1} = y_i + f(x_i, y_i)h + \frac{f'(x_i, y_i)}{2!} h^2 + \dots + \frac{f^{(n-1)}(x_i, y_i)}{n!} h^n + O(h^{n+1})$$

$$E_a \cong \frac{f'(x_i, y_i)}{2!} h^2 = O(h^2)$$

EULER

Local Truncation ERROR

- The error is reduced by 4 times if the step size is halved $\rightarrow O(h^2)$.
- In real problems, the derivatives used in the Taylor series are not easy to obtain.
- If the solution to the differential equation is *linear*, the method will provide error free predictions (2nd derivative is **zero** for a straight line).

Example: Euler's Method

Solve numerically : $\frac{dy}{dx} = -2x^3 + 12x^2 - 20x + 8.5$

From $x=0$ to $x=4$ with step size $h=0.5$

initial condition: $(x=0 ; y=1)$

Exact Solution: $y = -0.5x^4 + 4x^3 - 10x^2 + 8.5x + 1$

Numerical

Solution: $y_{i+1} = y_i + f(x_i, y_i)h$

$y(0.5) = y(0) + f(0, 1)0.5 = 1 + 8.5 * 0.5 = 5.25$

(true solution at $x=0.5$ is $y(0.5) = 3.22$ and $\epsilon_t = 63\%$)

$y(1) = y(0.5) + f(0.5, 5.25)0.5$
 $= 5.25 + [-2(0.5)^3 + 12(0.5)^2 - 20(0.5) + 8.5] * 0.5$
 $= 5.25 + 0.625 = 5.875$


(true solution at $x=1$ is $y(1) = 3$ and $\epsilon_t = 96\%$)

$y(1.5) = y(1) + f(1, 5.875)0.5 = 5.125$

....



X	y_{euler}	y_{true}	Error Global	Error Local
0	1	1	%	%
0.5	5.250	3.218	63.1	63.1
1.0	5.875	3.000	95.8	28
1.5	5.125	2.218	131.0	1.41
2.0	4.500	2.000	125.0	20.5



Improvements of Euler's method



- A fundamental source of error in Euler's method is that the derivative at the beginning of the interval is assumed to apply across the entire interval.
- Two simple modifications are available to circumvent this shortcoming:
 - **Heun's Method**
 - **The Midpoint** (or Improved Polygon) Method

Heun's method

- To improve the estimate of the slope, determine two derivatives for the interval:
 - At the initial point
 - At the end point
- The two derivatives are then averaged to obtain an improved estimate of the slope for the entire interval.

$$\textbf{Predictor} : y_{i+1}^0 = y_i + f(x_i, y_i)h$$

$$\textbf{Corrector} : y_{i+1} = y_i + \frac{f(x_i, y_i) + f(x_{i+1}, y_{i+1}^0)}{2}h$$



Heun's method (**improved**)

Original Huen's:

$$\mathbf{Predictor} : y_{i+1}^0 = y_i + f(x_i, y_i)h$$

$$\mathbf{Corrector} : y_{i+1} = y_i + \frac{f(x_i, y_i) + f(x_{i+1}, y_{i+1}^0)}{2} h$$

Note that the corrector can be iterated to improve the accuracy of y_{i+1} .

$$\mathbf{Predictor} : y_{i+1}^0 = y_i + f(x_i, y_i)h$$

$$\mathbf{Corrector} : y_{i+1}^j = y_i + \frac{f(x_i, y_i) + f(x_{i+1}, y_{i+1}^{j-1})}{2} h \quad j = 1, 2, \dots$$

However, it does not necessarily converge on the true answer but will converge on an estimate with a small error.


Example 25.5 from Textbook

The Midpoint (or Improved Polygon) Method


- Uses Euler's method to predict a value of y using the slope value at the midpoint of the interval:

$$y_{i+1/2} = y_i + f(x_i, y_i) \frac{h}{2}$$

$$y_{i+1} = y_i + f(x_{i+1/2}, y_{i+1/2})h$$



(a)



(b)

Runge-Kutta Methods (RK)

- Runge-Kutta methods achieve the accuracy of a Taylor series approach without requiring the calculation of higher derivatives.

$$y_{i+1} = y_i + \phi(x_i, y_i, h)h$$

$$\phi = a_1k_1 + a_2k_2 + \cdots + a_nk_n$$

Increment Function

a 's are constants

$$k_1 = f(x_i, y_i)$$

$$k_2 = f(x_i + p_1h, y_i + q_{11}k_1h)$$

p 's and q 's are constants

$$k_3 = f(x_i + p_3h, y_i + q_{21}k_1h + q_{22}k_2h)$$

⋮

$$k_n = f(x_i + p_{n-1}h, y_i + q_{n-1}k_1h + q_{n-1,2}k_2h + \cdots + q_{n-1,n-1}k_{n-1}h)$$

Runge-Kutta Methods (cont.)

- Various types of RK methods can be devised by employing different number of terms in the increment function as specified by n .
- **First order** RK method with $n=1$ and $a_j=1$ is in fact **Euler's method**.

$$y_{i+1} = y_i + \phi(x_i, y_i, h)h$$

$$\phi = a_1k_1 + a_2k_2 + \cdots + a_nk_n$$

$$k_1 = f(x_i, y_i)$$

choose $n=1$ and $a_1=1$, we obtain

$$y_{i+1} = y_i + f(x_i, y_i)h \quad \text{(Euler's Method)}$$

Runge-Kutta Methods (cont.)

Second-order Runge-Kutta Methods:

$$y_{i+1} = y_i + (a_1k_1 + a_2k_2)h$$

$$k_1 = f(x_i, y_i) \quad k_2 = f(x_i + p_1h, y_i + q_{11}k_1h)$$

- Values of \mathbf{a}_1 , \mathbf{a}_2 , \mathbf{p}_1 , and \mathbf{q}_{11} are evaluated by setting the above equation equal to a *Taylor series expansion* to the second order term. This way, three equations can be derived to evaluate the four unknown constants (See **Box 25.1** for this derivation).

A value is assumed for one of the unknowns to solve for the other three.

$$a_1 + a_2 = 1$$

$$a_2p_1 = \frac{1}{2}$$

$$a_2q_{11} = \frac{1}{2}$$

$$y_{i+1} = y_i + (a_1k_1 + a_2k_2)h$$

$$k_1 = f(x_i, y_i)$$

$$k_2 = f(x_i + p_1h, y_i + q_{11}k_1h)$$

$$a_1 + a_2 = 1$$

$$a_2p_1 = \frac{1}{2}$$

$$a_2q_{11} = \frac{1}{2}$$

- Because we can choose an infinite number of values for a_2 , there are an infinite number of second-order RK methods.

Three of the most commonly used methods are:

– **Huen's Method** with a Single Corrector ($a_2=1/2$)

– **The Midpoint Method** ($a_2=1$)

– **Ralston's Method** ($a_2=2/3$)

Huen's Method ($a_2= 1/2$) → $a_1= 1/2$ $p_1= 1$ $q_{11}= 1$

$$y_{i+1} = y_i + (a_1k_1 + a_2k_2)h = y_i + \left(\frac{1}{2}k_1 + \frac{1}{2}k_2\right)h$$

$$k_1 = f(x_i, y_i)$$

$$k_2 = f(x_i + h, y_i + k_1h)$$

$$y_{i+1} = y_i + (a_1 k_1 + a_2 k_2)h$$

$$k_1 = f(x_i, y_i)$$

$$k_2 = f(x_i + p_1 h, y_i + q_{11} k_1 h)$$

$$a_1 + a_2 = 1$$

$$a_2 p_1 = \frac{1}{2}$$

$$a_2 q_{11} = \frac{1}{2}$$

The Midpoint Method ($a_2 = 1$)



$$a_1 = 0$$

$$p_1 = 1/2$$

$$q_{11} = 1/2$$

$$y_{i+1} = y_i + (a_1 k_1 + a_2 k_2)h = y_i + (k_2)h$$

$$k_1 = f(x_i, y_i) \quad k_2 = f\left(x_i + \frac{1}{2}h, y_i + \frac{1}{2}k_1 h\right)$$

$$y_{i+1} = y_i + (k_2)h = y_i + f\left(x_i + \frac{h}{2}, y_i + \frac{h}{2}f(x_i, y_i)\right)h$$

- Three most commonly used methods:


- **Huen Method** with a Single Corrector ($a_2=1/2$)

- **The Midpoint Method** ($a_2=1$)

- **Ralston's Method** ($a_2=2/3$)

Ralston's Method ($a_2=2/3$)

Comparison of Various Second-Order RK Methods



$$y_{i+1} = y_i + (a_1 k_1 + a_2 k_2)h = y_i + \left(\frac{1}{3}k_1 + \frac{2}{3}k_2\right)h$$

$$k_1 = f(x_i, y_i)$$

$$k_2 = f\left(x_i + \frac{3}{4}h, y_i + \frac{3}{4}k_1 h\right)$$

Systems of Equations

- Many practical problems in engineering and science require the solution of a system of simultaneous ordinary differential equations (ODEs) rather than a single equation:

$$\begin{aligned}\frac{dy_1}{dx} &= f_1(x, y_1, y_2, \dots, y_n) \\ \frac{dy_2}{dx} &= f_2(x, y_1, y_2, \dots, y_n) \\ &\vdots \\ \frac{dy_n}{dx} &= f_n(x, y_1, y_2, \dots, y_n)\end{aligned}$$

- Solution requires that n initial conditions be known at the starting value of x .
i.e. $(x_0, y_1(x_0), y_2(x_0), \dots, y_n(x_0))$
- At iteration i , n values $(y_1(x_i), y_2(x_i), \dots, y_n(x_i))$ are computed.