

BREAST CANCER RESEARCHES IN SAUDI ARABIA

Prepared by
DR. SAMIA AL-AMOUDI



جادة الابداع
JADDAT AL-EBDAA

055 4679687
tojadat@yahoo.com

MEMBERS OF SHEIKH MOHAMMED HUSSEIN AL-AMOUDI SCIENTIFIC CHAIR FOR BREAST CANCER RESEARCHES



Dr. Samia Al-Amoudi

Chairwomen of the Scientific Chair for Breast Cancer researches.
CEO and Board member of sheikh Mohammed Hussein Al-Amoudi Center of Excellence in Breast Cancer.
Breast Cancer Survivor, Global Advocate Susan G Komen for the Cure, USA.
King Abdulaziz University, Jeddah, Saudi Arabia.



Prof. Mahmoud Shaheen Al-Ahwal

Professor of Internal Medicine and Medical Oncology.
Dean, Faculty of Medicine. KAU, Jeddah, Saudi Arabia.
Member of the Scientific Chair for Breast Cancer researches.
Board member of sheikh Mohammed Hussein Al-Amoudi Center of Excellence in Breast Cancer.
King Abdulaziz University, Jeddah, Saudi Arabia.



Dr. Wafa Sait

Consultant obstetrician gynecologist and operative endoscopy.
Department of obstetric and Gynecology.
Member of the Scientific Chair for Breast Cancer researches.
Board member of sheikh Mohammed Hussein Al-Amoudi Center of Excellence in Breast Cancer.
King Abdulaziz University , Jeddah, Saudi Arabia.



Dr. Zuhoor K. Al-Ghaithy

Associate Professor and Consultant general surgery oncoplastic breast surgery
Department of Surgery.
Member of the Scientific Chair for Breast Cancer researches.
Board member of sheikh Mohammed Hussein Al-Amoudi Center of Excellence in
Breast Cancer.
King Abdulaziz University , Jeddah, Saudi Arabia.



Dr. Yasir Abdul Aziz Bahader

Professor of Radiation Oncology.
Chairman of Radiology Department.
Member of the Scientific Chair for Breast Cancer researches.
King Abdulaziz University , Jeddah, Saudi Arabia.



Dr. Bassem Salama El-Deek

Associate Professor Community Medicine.
Joint master of health profession education (JMHPE) Maastricht.
KAU FOM, Medical education department .
Member of the Scientific Chair for Breast Cancer researches.
King Abdulaziz University , Jeddah, Saudi Arabia.



Omar Alzyout

Research Assistant at Sheikh Mohammed Hussein Al-Amoudi Scientific Chair for
Breast Cancer.
Master degree in Social Anthropology/ Yarmouk University - Irbid - Jordan.
Bachelor degree in Sociology/ The University of Jordan - Amman - Jordan.
King Abdulazeez University - Jeddah - Saudi Arabia.

5	EDITORIAL Professor Mahmoud Shaheen Al-Ahwal
6	ASSESSMENT OF THE QUALITY OF BREAST CANCER CARE: A SINGLE INSTITUTIONAL STUDY FROM SAUDI ARABIA Shad Salim Akhtar and Hisham Mohammed Nadrah
13	KNOWLEDGE OF BREAST CANCER AND ITS RISK AND PROTECTIVE FACTORS AMONG WOMEN IN RIYADH Awatif Ali Alam
20	THE KNOWLEDGE OF BREAST CANCER AMONG YOUNG SAUDI FEMALES Wafa A. Sait, Samia M. Al-Amoudi, Dalal A.Tawtai, Hassan S. Abduljabbar
25	FNA OF BREAST FIBROADENOMA: OBSERVER VARIABILITY AND REVIEW OF CYTOMORPHOLOGY WITH CYTOHISTOLOGICAL CORRELATION S. M. Kollur and I. A. El Hag
32	DIETARY FAT AND BREAST CANCER IN SAUDI ARABIA: A CASE-CONTROL STUDY A. Alothaimeen, A. Ezzat, G. Mohamed, T. Muammar and A. Al-Madouj
39	PATTERN OF CANCER IN SAUDI ARABS REFERRED TO KING FAISAL SPECIALIST HOSPITAL Salah M. El-Akkad, Magid H. Amer, Gregory S. Lin, Rajeh S. Sabbah, and John T. Godwin
47	PATTERN OF CANCER AT ASIR CENTRAL HOSPITAL, ABHA, SAUDI ARABIA Abdur Rauf Khan, Nazir K. Hussain, Abdulaziz Al-Saigh, Tarek Malatani, Anwar A. Sheikha
52	CANCER IN AL-QASSIM, SAUDI ARABIA: A RETROSPECTIVE STUDY (1987-1995) Shad Salim Akhtar, Lolita Malig Reyes
60	THE SPECTRUM OF BREAST DISEASES IN SAUDI ARAB FEMALES: A 26 YEAR PATHOLOGICAL SURVEY AT DHAHRAN HEALTH CENTER Shad Samir S. Amr, Abdul Rahman M. Sa'di, Fazal Ilahi, Salwa S. Sheikh



EDITORIAL

Dear colleagues,

Breast cancer is the most common cancer among Saudi males and females according to the Cancer Incidence report from the National Cancer Registry of the kingdom of Saudi Arabia. Despite the low Cancer Incidence in the Kingdom of Saudi Arabia, because of the high percentage of young population, we must be ready to face the expected increase in cancer burden attributed to growth, aging of population and awareness of our society which will lead to tremendous increase in detecting breast cancer early and hence, significant reduction in mortality. Research promotion in general is one of the important steps in identifying breast cancer epidemiology, causation, the need for breast cancer screening and health care planning.

This publication is one step to promote awareness and education among physicians, nurses and health care providers in the KSA and hence more research in breast cancer. In this occasion, and on behalf of Faculty of Medicine, King Abdulaziz University, I would like to thank Dr. Samia Al-Amoudi – CEO of Sheikh Mohammed Hussien Al-Amoudi Center of Excellence in Breast Cancer, for her great effort at national and international levels, leading the society towards a new era in women's health and especially in breast cancer.

Prof. Mahmoud Shaheen Al-Ahwal

Professor of Internal Medicine and Medical Oncology

Dean, Faculty of Medicine. KAU, Jeddah, Saudi Arabia.

Member of Sheikh Mohammed Hussein Al-Amoudi Scientific Chair for Breast Cancer

ASSESSMENT OF THE QUALITY OF BREAST CANCER CARE: A SINGLE INSTITUTIONAL STUDY FROM SAUDI ARABIA*

SHAD SALIM AKHTAR AND HISHAM MOHAMMED NADRAH
King Fahd Specialist Hospital, Prince Faisal Oncology Center, Buraidah, Al-Qassim, Saudi Arabia

ABSTRACT

Objective: To evaluate the quality of operable breast cancer care in a tertiary care institution.

Design: A retrospective analysis of all breast cancer patients seen in our institution between 1995 and 2000. Data were abstracted from the charts of these patients. Indicators were based on an international consensus conference and other publications.

Setting: A tertiary care health care institution.

Main measures: We evaluated the charts and calculated the percentage for which the internationally accepted quality care indicators were followed during the continuum of care. We also reviewed the histopathological reports to evaluate conformation with the accepted indicators.

Results: Charts of 75 patients (four exclusions, three metastatic, and one male), diagnosed to have breast cancer during the study period were reviewed. Only 28 (37%) patients had triple assessment before a definitive surgical procedure. Pre-operative staging including a CT and bone scan was performed in 58 (77.3%). Among the 50 patients who had definite surgical intervention, the majority had mastectomy (44/50, 88%) whereas axillary dissection was performed in 46 (46/50, 92%). Estrogen and progesterone receptor status was reported in only four (4/50, 8%) and the exact tumor size in 24 (24/50, 48%) of the histopatho-

* International Journal for Quality in Health Care 2005, Vol. 17, No. 4: pp.301-305

logical reports. Adjuvant chemotherapy was used in accordance with the international standards but radiotherapy was under-utilized.

Conclusion: Our study demonstrated that the quality of breast cancer care in this institution was below the accepted international standards. This study may be used to make interventions for improvement of quality in similar institutions all over the kingdom.

Keywords: assessment of care, breast cancer, care quality, health care quality, quality assessment

Defined as 'the degree to which health services for individuals and populations increase the likelihood of the desired outcomes and are consistent with current professional knowledge'; quality of care is a matter of concern both for the care providers and the health care recipients [1-3]. Process quality, which refers to what health care workers do, and how well they do it, both technically as well as interpersonally, is commonly used to assess the quality of care. Particularly the quality of the technical

process is measured, which refers to whether the right choices are made in diagnosing and treating the patient and whether care is provided in an effective and skillful manner. Obviously the best process measures are those for which there is evidence from research that a better process leads to an improved outcome [4]. Breast cancer is the most common cancer diagnosed in the world among women. During all phases of continuum of breast cancer care, an association between the process and outcome is supported by extensive scientific literature including many randomized trials, meta-analyses, and international guidelines [5-8]. It therefore provides an excellent opportunity to assess the quality of care among this group of oncological patients. For this reason, even though the age-standardized rate for breast cancer is quite low (4.73/100 000) in the Al-Qassim region of Saudi Arabia [9], we conducted a retrospective study to assess the quality of care offered to these patients at the King Fahd Specialist Hospital and its Prince Faisal Oncology Center.

PATIENTS AND METHODS

According to the 1999 census, the population of

Table 1: Indicators used for the assessment of quality of care in breast cancer patients 1995-2000

Parameter	Expert target (%)	Reference
Diagnosis		
Pre-operative mammography	~100	[21]
Pre-operative biopsy ¹	>95	[11]
Non-diagnostic fine needle biopsy	<10	[13]
Triple assessment	~100	[16,22]
Pre-operative staging	>94	[12]
Surgical treatment		
Breast-conserving treatment	~50	[11]
Axillary dissection	>90	[11]
Histopathological reporting		
Tumor size	>99	[17,18,23,29]
Grade	>99	[19,23]
ER/PR status	>99	[17-19,23]
Number of nodes	>99	[17,18,23]
Adjuvant chemotherapy		
Pre-menopausal	>90	[11]
Post-menopausal	~50	[11]

ER/PR, estrogen receptor/progesterone receptor.

¹Biopsy could be aspiration cytology, core biopsy, or excisional biopsy.

the Al-Qassim region was 933 146, of which 427 507 were females [10]. The King Fahd Specialist Hospital, run by the Ministry of Health, is the main referral hospital of the region. To streamline the management of cancer patients the Prince Faisal Oncology Center was commissioned within this institution in 1998. The King Fahd Specialist Hospital and its Prince Faisal Oncology Center at Buraidah receive the majority of patients suspected or diagnosed to have malignant disease, from all over Al-Qassim region. The latter also maintains a hospital-based cancer registry for the region. We identified breast cancer patients from our medical record indexing system and the database of Prince Faisal Oncology Center. Data abstracted from the charts of 78 breast cancer patients, seen in our institution between 1995 and 2000, were copied into a blank proforma.

To study the quality of care in this group of patients, indicators proposed by the National Cancer Policy Board [4] and Hillner et al.[11] were used. In addition, standards of practice outlined in various guidelines and reviews were utilized to assess other parameters of breast cancer care (Table 1) [8,12-15].

The quality of fine needle aspiration cytology was assessed by dividing our patients into two groups: those whose diagnosis was established by this procedure and others who needed a subsequent core or excision biopsy before a definitive surgical procedure due to non-diagnostic aspiration cytology.

The accuracy of clinical examination, mammography, and fine needle aspiration cytology increases when they are used together; a procedure called triple assessment or triple diagnosis.

When the results of all the components of triple assessment indicate a benign lesion, cancer will be found in less than 0.5% of cases. On the other hand if all three indicate cancer the diagnosis is likely to be confirmed in more than 99% of cases [14,18]. The size of the tumor was abstracted from the pathological, surgical, or clinical record of the patient, whichever was available. When all three were documented pathological size took precedence. Pre-operative staging was considered complete if the patient had a chest X-ray, liver function tests, abdominal ultrasound, computerized tomography scan, and a bone scan.

Clinical and pathological staging information was additively utilized to identify the stage of disease according to the American Joint Committee for Staging (AJCC) classification [23].

We reviewed the histopathological reports of all the biopsies as well as post-operative specimens

of those patients who had a definitive surgical procedure. Table 1 shows the criteria used to evaluate the quality of histopathological reporting.

Table 2: Characteristics of breast cancer patients 1995-2000

Parameter	No. (%)
Age at diagnosis (years)	
Mean (range)	46 ± 14.7 (23-100)
≤40	32/75 (42.7)
41-50	24/75 (32)
>51	19/75 (25.3)
Pre-menopausal	55/75 (73.3)
Post-menopausal	20/75 (26.7)
Symptoms at presentation	
Lump	74/75 (98.7)
Nipple discharge	1/75 (1.3)
Tumor size distribution ¹	
T1 (<2 cm)	3/75 (4)
T2 (2-5 cm)	25/75 (33.3)
T3 (>5 cm)	23/75 (30.7)
T4 ²	17/75 (22.7)
Tx (unknown)	7/75 (9.3)
Stage distribution	
II	30/75 (40)
III	22/75 (29.3)
Unknown	23/75 (30.7)

Stage II, T0N1M0, T1N1M0, T2N0M0, T2N1M0, T3N0M0; stage III, any TN2M0, T3N1M0 (N, lymph node involvement; M0, no distant metastasis).

¹Based on clinical, operative, or final pathological evaluation.

²T4, any T with direct extension to chest wall and/or skin.

RESULTS

During the 5-year period from 1995, 78 female breast cancer patients were seen in our institution. A lone male was excluded. As shown in Table 2 the mean age at diagnosis of our patients was 46 ± 14.7 years (range 23-110 years). Three of these patients were detected to have metastatic disease at presentation and were excluded from further analysis. Pre-operative bilateral mammogram was performed in 39 (39/75,52%) and fine needle aspiration cytology in 46 patients (46/75, 61.3%) (Table 3). Only 28 patients (28/75,37.3%) had triple assessment before a definitive surgical procedure. A core biopsy (27/75, 36%) or excisional biopsy

(24/75, 32%), was performed in 51 (51/75, 68%) patients and 22 patients (22/46, 47.8%) had this procedure after a non-diagnostic aspiration biopsy. Exact tumor size, either clinically, peri-operatively, or pathologically, had been documented in 68 patients (68/75, 90.6%). The majority of these patients had large tumors (T3 + T4 = 40/68, 58.8%). Fifty-two of the 58 patients (58/75, 77.3%) who had complete pre-operative staging, had adequate data available for staging according to the AJCC. The majority had advanced stage disease (*Table 2*).

Fifty (50/75, 66.6%) patients underwent a definitive surgical procedure in our institution, 44 (44/75, 58.7%) had mastectomy and the others (6/75, 8%) had only lumpectomy. The remainder (25/75, 33.3%) either refused surgery here or demanded referral elsewhere. No follow-up records were available for these patients. *Table 3* also shows the indicators as recorded in the histopathological reports of these patients.

Adjuvant hormone therapy was used in 24 pre-menopausal (24/38, 63.2%) and 11 post-menopausal (11/12, 91.6%) patients. Rational use of hormone therapy could not be assessed because estrogen and progesterone receptor status was not available in the majority of these patients. Twenty-five patients (20/38, 52.6% pre-menopausal; 5/12, 41.6% post-menopausal) received adjuvant radiotherapy.

DISCUSSION

The large number of studies that have been conducted to assess the process of breast cancer care in different settings, makes this extensive literature difficult to summarize. Our study, a single institutional study, assessed the quality of care during a period of time using the data abstracted from the medical records of these patients. Like a recently published series, this study albeit small, confirms the feasibility of assessing the quality of cancer care by abstracting the data from the medical records of patients [3]. Similarly, the American Society of Clinical Oncology-sponsored National Initiative for Cancer Care Quality is currently evaluating the feasibility of getting information from the medical records of patients to assess the quality of care on a large scale [24].

This study highlights some of the unique epidemiological features of breast cancer seen in this region; younger age, large tumor size, and advanced stage at presentation. In contrast to data from the West, where more than half of breast cancer patients are above 50 years of age, the majority of our patients were younger than that (*Table 2*). This age

distribution has been reported previously from this area [25]. Racial difference in stage at presentation and aggressive disease in younger patients have been described in the literature [26,27].

A younger age at presentation may be one of the reasons for the advanced stage of disease seen in our patients. However, lack of awareness regarding breast cancer and delay in seeking medical advice may be additional factors contributing to this late presentation [28].

Ideally, every breast cancer patient should have a bilateral mammogram before the definitive surgical procedure for primary disease [16]. Similarly, triple assessment is an important part of the evaluation of any breast lump [17]. Both mammography and as a result, the triple assessment, were under-utilized in our patients. Among other factors, temporary non-availability of mammography in our institution, at the time of referral of some of these patients, could have contributed to the low rate of pre-operative bilateral mammography seen in our patients.

Although an operator-dependent procedure, the diagnostic value of fine needle aspiration cytology is reported to be ~90% with a false-negative rate ranging between 0.4 and 35% [4,13].

As most of the clinicians would themselves carry out this procedure in this institution, the high rate of non-diagnostic fine needle biopsy seen by us may indicate an initial learning curve.

In order to plan adequate treatment, especially breast-conserving surgery, every breast cancer patient needs evaluation by a team consisting of a surgeon, medical oncologist, radiation oncologist, pathologist, and a radiologist. Non-availability of radiotherapy and barriers to patient follow-up may possibly explain why only a small proportion of patients undergoing surgery in our institution had lumpectomies. Nevertheless, patient preference and physician's choice are two of the other factors that may override the medical criteria and affect the rate of breast-conserving surgery in an institution [29].

In conformity with international standards, all but four patients undergoing definitive surgery here had axillary dissection.

The pathological report is a critical link between pathologist and clinician. Deficits in pathological reports have, therefore, been the target of many quality improvement projects.

In our study the histopathological reporting did not meet the expected international standards. Although a low rate of reporting of estrogen and progesterone receptor status could be due to non-availability of such marker studies until 1999 in our

institution, other aspects of histopathological reporting were also below the acceptable standard (Table 3). The number of incomplete breast cancer pathology reports was recently reduced as reported by Hammond and Flinner [22] from 57/356 (16%) to 2/190 (1.1%) by instituting a template for reporting.

The rate of utilization of adjuvant chemotherapy both in pre-menopausal and post-menopausal patients met the expected target [11]. Because chemotherapy can be made easily available there is a high rate of concordance. Adjuvant radiotherapy is an integral part of the management of breast cancer, particularly in patients with large tumors and many positive nodes [30]. Most of our patients had large primary tumors with axillary lymph node involvement and were, therefore, candidates for local adjuvant radiotherapy. Again, perhaps due to non-availability of radiotherapy on site, it was under-utilized in this patient population.

CONCLUSION

This small retrospective analysis shows that quality of breast cancer care in this institution falls well below the accepted international standards, possibly due to non-availability of some of the facilities on site and the absence of local guidelines. Additionally, a low volume of breast cancer patients in this institution may have contributed. Since the compilation of this report a clinic for fine needle aspiration cytology has been established and local guidelines have been formulated including a template for histopathological reporting. Collaboration with an institution for radiotherapy facilities has been established.

It would be interesting to review the status of practice in future and compare that with the present study.

Table 3: Compliance with the guidelines as observed in breast cancer patients 1995-2000

Parameter	No. having the procedure (%)
Diagnosis	
Pre-operative mammography	39/75 (52)
Pre-operative biopsy	
Fine needle biopsy	46/75 (61.3)
Core biopsy	27/75 (36)
Excisional biopsy	24/75 (32)
Non-diagnostic fine needle biopsy	22/46 (47.8)
Triple assessment	28/75 (37.3)
Pre-operative staging	58/75 (77.3)
Surgical treatment	
Breast-conserving surgery	2/22 (9.1)
Axillary dissection	46/50 (92)
Histopathological reporting	
Tumor size	24/50 (48)
Grade	36/50 (72)
ER/PR status	4/50 (8)
Total number of nodes removed	39/50 (78)
Adjuvant chemotherapy	
Pre-menopausal	34/38 (89.5)
Post-menopausal (58.3)	7/12

FNAC, fine needle aspiration cytology; ER/PR, estrogen receptor/progesterone receptor.

REFERENCES

1. Lohr KN, ed., Medicare: A Strategy for Quality Assurance. Washington DC: National Academy Press, 1990.
2. Chassin MR, Gavin RW. The urgent need to improve health care quality. Institute of Medicine National Round Table on Health Care Quality. JAMA 1998; 280: 1000–1005.
3. McGlynn EA, Asch SM, Adams J et al. The quality of health care delivered to adults in the United States. N Engl J Med 2003; 348:2635–2645.
4. Hewitt ME, Simone JV, National Cancer Policy Board, Institute of Medicine and National Research Council. Ensuring Quality Cancer Care. Washington, DC: National Academy Press, 1999.
5. Eastman P. NCI adopts new mammography screening guidelines for women. J Natl Cancer Inst 1997; 89: 538–539.
6. Early Breast Cancer Trialists' Collaborative Group. Systemic treatment of early breast cancer by hormonal, cytotoxic, or immune therapy. 133 randomized trials involving 31,000 recurrences and 24,000 deaths among 75,000 women. Lancet 1992; 339: 1–71.
7. Early Breast Cancer Trialists' Collaborative Group. Tamoxifen for early breast cancer: an overview of the randomized trials. Lancet 1998; 351: 1451–1467.
8. Goldhirsch A, Glick J H, Gelbr RD et al. Meeting highlights: International Consensus Panel on the treatment of primary breast cancer. J Natl Cancer Inst 1998; 90: 1601–1608.
9. National Cancer Registry. Cancer Incidence Report Saudi Arabia 1994–1996. Riyadh: Ministry of Health, 1999.
10. Central Department of Statistics, Kingdom of Saudi Arabia: <http://www.planning.gov.sa/statistic> Accessed 16 July 2004.
11. Hillner BE, Mc Donald MK, Penberthy L et al. Measuring standards of care for early breast cancer in an insured population. J Clin Oncol 1997; 15: 1401–1408.
12. Fleming ID, Philips JL, Menck HR, Murphy GP, Winchester DP. The National Cancer Data Base report on recent hospital cancer program progress toward complete American Joint Committee on Cancer/ TNM staging. Cancer 1997; 80: 2305–2310.
13. Kline TS. Survey of aspiration biopsy cytology of the breast. Diagn Cytopathol 1991; 7: 98–105.
14. Donegan WL. Evaluation of a palpable breast mass. N Engl J Med 1992; 327: 937–942.
15. Mandelblatt JS, Ganz PA, Kahn KL. Proposed agenda for the measurement of quality-of-care outcomes in oncology practice. J Clin Oncol 1999; 17: 2614–2622.
16. Grilli R, Apolone G, Marsoni S et al. The impact of patient management guidelines on the care of breast, colorectal and ovarian cancer patients in Italy. Med Care 1991; 29: 50–63.
17. The Steering Committee on Clinical Practice Guidelines for the Care and Treatment of Breast Cancer. The palpable breast lump: information and recommendations to assist decision making when a breast lump is detected. Can Med Assoc J 1998; 158: s3–s8.
18. Kaufman Z, Shpitz B, Shapiro M et al. Triple approach in the diagnosis of dominant breast masses. Combined physical examination, mammography and fine needle aspiration. J Surg Oncol 1994; 56: 254–257.
19. Hand R, Sener S, Imperato J et al. Hospital variables associated with quality of care for breast cancer patients. JAMA 1991; 266:3429–3432.

20. Ostein RT. Breast cancer. In Steele GD, Winchester DP, Menck HR, Murphy GP, eds, National Cancer Data Base. Atlanta,GA: American Cancer Society, 1992: 29–38.
21. Nakleh RE, Jones B, Zarbo RJ. Mammographically directed breast biopsies. A college of American Pathologists Q-probes study of clinical physician expectations and of specimen handling and reporting characteristics in 434 institutions. *Arch Pathol Lab Med* 1997; 121: 11–18.
22. Hammond EH, Flinner RL. Clinically relevant breast cancer reporting. Using process measures to improve anatomic pathology reporting. *Arch Pathol Lab Med* 1997; 1: 1171–1175.
23. Beahrs OH, Henson DE, Hutter RVP, Kennedy BJ, eds, American Joint Committee on Cancer: Manual for Staging of Cancer, 4th edn. Philadelphia: JB Lippincott Co. 1992.
24. Malin JL, Schuster MA, Kahn KA, Brook RH. Quality of breast cancer care: What do we know? *J Clin Oncol* 2002; 20: 4381–4393.
25. Akhtar SS, Reyes LM. Cancer in Al-Qassim, Saudi Arabia: A retrospective study (1987–1995). *Ann Saudi Med* 1997; 17: 595–560.
26. de la Rochefordiere A, Asselain B, Campana F et al. Age as a prognostic factor in pre-menopausal breast carcinoma. *Lancet* 1993; 341: 1039–1043.
27. Jones BA, Kasl SV, Curnen MGM, Owens PH, Dubrow R. Can mammography screening explain the race difference in stage at diagnosis of breast cancer? *Cancer* 1995; 75: 2103–2113.
28. Akhtar SS, Filani T, Gadella M, Amri A. Beliefs and attitudes about breast self-examination in Al-Qassim region of Saudi Arabia. A study of women attending primary health care centers. *Saudi Med J* 1995; 16: 493–497.
29. Hotobagyi GN. Treatment of breast cancer. *N Engl J Med* 1998; 339: 974–984.
30. Overgaard M. Radiotherapy as a part of a multidisciplinary treatment strategy in early breast cancer. *Eur J Cancer* 2001; 37: S33–S43.

KNOWLEDGE OF BREAST CANCER AND ITS RISK AND PROTECTIVE FACTORS AMONG WOMEN IN RIYADH*

AWATIF ALI ALAM

Department of Family and Community Medicine, King Saud University, Riyadh, Saudi Arabia

ABSTRACT

Background: We conducted this study to assess knowledge of breast cancer and sources of information about breast cancer among women in Riyadh. We also analyzed whether associations existed between demographic variables, knowledge of breast cancer, and the practice of breast self-examination and use of mammography screening.

Methods: Women interested in participating in this community-based descriptive study provided data by completing a pre-tested structured questionnaire.

Results: Of 864 participating women, 84% were Saudi, 45% were married and 67.8% had a university level education. Eighty percent were between the ages of 20 to 50 years. Knowledge of breast self examination (BSE) was high; 82% (95% confidence intervals [CI], 79.2%-84.4%) knew about BSE, while 61% (95% CI: 57.9%-64.5%) knew about mammography, but only 41.2% (95% CI, 37.9%-44.5%) had performed BSE and 18.2% (95%CI, 15.5%-20.8%) had had mammography screening. Knowledge of breast cancer, risk factors and protective factors for breast cancer was moderate. There was a statistically significant association between the demographic characteristics (marital status, educational status and family history of breast cancer) and knowledge and practice of BSE and mammography.

Conclusion: Though it has limitations, this study revealed an imbalance between the knowledge and practice

* Annual of Saudi Medicine 2006, 26(4):272-7

of BSE among women. It also showed that there is only moderate knowledge of risk and protective factors for breast cancer and that knowledge and practice of BSE and mammograms vary according to marital and educational status. Hence, frequent community-based awareness programs are needed so that all women can know and practice BSE, which in turn helps to prevent breast cancer.

Breast cancer continues to be a major cause of morbidity and mortality throughout the world. With 1 million new cases in the world each year, breast cancer is the commonest cancer in women, comprising 18% of all female cancers. Of every 1000 women 50 years of age, two will have recently had breast cancer diagnosed and about 15 will have had a diagnosis made before the age of 50 years, giving a prevalence of breast cancer of nearly 2%.^[1] The incidence of breast cancer increases up to the age of 80 years, plateaus between the ages of 80 to 85 years, and then declines. However, the measured decline after the age of 85 years is difficult to interpret and may reflect the inadequacy of the epidemiologic data. The prevalence of breast cancer in recent years has prompted women to seek medical advice randomly with minimal breast symptoms, but only a small number of women are aware of the the society, with childbearing extending practically over the entire reproductive period of life. Due to the conservative nature of the society, many females will refrain from seeking medical advice out of shyness until their disease becomes far advanced, particularly in cases of carcinoma of the breast. Often Breast cancer engenders an exceptional level of fear among women, most probably because of its external location on the body, with all of the obvious cosmetic and psychosocial implications, coupled with proper methods of conducting breast self examination (BSE) or are aware of the importance of radiological screening for breast cancer.^[2] Breast cancer is known to be the most common malignancy among women internationally and the Kingdom of Saudi Arabia is no exception.^[3] Certain social factors and demographic findings contribute to the pattern of disease of the breast encountered in Saudi Arabia.^[4] Saudi Arab females tend to marry at a young age, according to the traditional conservative values of they fear the treatment more than the disease itself.^[4] the major concern with all types of cancer, namely loss of life. This places an even greater burden upon the healthcare system to convince such patients to undergo screening for not only will earlier disease be more curable, it will more likely be treatable without loss of the breast. Paramount to early detection in

an organized screening program is physician examination, at appropriate intervals, coupled with patient self-examination where this is acceptable,^[5] and the efforts of the nursing staff in various clinics to encourage females to seek medical help if they feel a breast lump. However, such efforts are more successful among younger females who are using the available facilities, such as the breast clinic and most importantly, mammography.^[4] In the United States, factors considered statistically "protective" against breast cancer include early child bearing, multiparity and breastfeeding. All of these factors are commonly present among Saudi women, yet not only do they get breast cancer anyway, but frequently at an age more than a decade younger than their American counterparts.^[5]

In the Cancer Incidence Report for Saudi Arabia, published in May 1999, breast cancer accounted for 19.1% of cancers among females whose mean age at diagnosis was 48.3 years. Diagnosis of in situ carcinoma was almost nil, while infiltrating ductal carcinoma was diagnosed in 76.1 %.^[6] In a study of 130 women with invasive breast cancer in Saudi Arabia, 82% were under 50 years of age and 16% were under 30 years of age. This suggests that the lower age limit for breast cancer screening, when un-dertaken in Saudi Arabia, should be younger than the European recommendation, which states that breast screening could start at age 40, but the most beneficial age is 50 years and older.^[7] El-Harith et al concluded that BRCA1 and BRCA2 mutations are likely to contribute to the pathogenesis of familial breast cancer in female patients from Saudi Arabia.^[8] Breast and lung cancer share the second position as leading causes of death from cancer among Saudi females, with each being responsible for 10% of the total deaths from cancer.^[3] Delays in diagnosis and management are associated with advancing disease and consequently poorer results with any appropriate treatment.^[9] The incidence of breast carcinoma is increasing in the Kingdom. According to the 2001 Tumor Registry Annual Report published by King Faisal Specialist Hospital and Research Center, 348 cases were seen in 2001 compared with 131 cases seen in 1985.^[10] The incidence rate of cancer in Saudi Arabia is predicted to continue rising, partly due to rapid changes in lifestyle, the increasing consumption of cigarettes, the high caloric and fatty diet intake observed during the last decade, and partly due to the natural aging of this young Saudi population in the years to come.^[11]

For mammography to be useful, it must be available to the population of women at risk. There must

be sufficient numbers of well-trained, experienced mammographers to interpret the films, and there must be methodology available to define and diagnose mammographically detected lesions. Physicians must also emphasize the critical application of screening mammography to the early detection and treatment of breast cancer. Therefore, the care of patients should be enhanced as should the education of those in training in this area. It provides a richer professional environment for all involved in the care of breast cancer patients, and provides a fertile ground for exchange of ideas for research. Ongoing research projects are enhanced by this exchange, and ideas for new projects have been spawned.^[5]

Generally, classical medical care for patients with breast and ovarian cancers is easily accessible in most of the major medical centers of Saudi Arabia. The diagnostic facilities available in this country are quite adequate and they include, other than mammography, routine medical examination, fine needle aspiration and other advanced histopathological techniques. However, genetic testing for assessment of predisposing to breast cancer is presently not offered.^[3] The spectrum of treatment services offered includes lumpectomy, mastectomy with or without breast reconstitution, chemotherapy and/or radiotherapy.

The greatest advantage to the women of the Kingdom will be the development of more widespread screening and educational programs. Currently, lectures in a major hospital in Riyadh that foster public awareness of breast cancer are given every two months and certified nurses provide workshops on breast-examination.^[11] Breast cancer awareness programs (BCAP) are meant to help in the prevention and early detection of breast cancer. This study was carried out with the objectives of (i) assessing the knowledge of breast cancer and its associated factors (risk and protective) among the women of Riyadh city, (ii) assessing the source of knowledge of breast cancer, (iii) observing any association between family history of breast cancer, marital status and educational level of study participants versus their knowledge and practice of breast self examination and mammography.

MATERIALS AND METHODS

A community-based descriptive study was conducted on a sample of 864 adult women during a health education campaign held in October 2003 in the Ladies Kingdom Mall in Riyadh. Women willing to participate, regardless of their age or educational level were included in the study.

A questionnaire was distributed to participants that asked about sociodemographic factors such as age, nationality, occupation, marital status, and level of education. It also included questions about the changes that occur in relation to breast cancer, risk factors that increase susceptibility, management practices for breast cancer, protective factors against breast cancer and sources of knowledge. The data was analyzed using SPSS-PC statistical software. For descriptive statistics, 95% confidence intervals were used to show the precision of the estimates. The chi-square test was used to test for association.

RESULTS

The sample consisted of 864 women of different ages, with 80% between the ages of 20 to 50 years. Details of the age structure of participants and other sociodemographic characteristics are shown in *Table 1*. Saudi nationals represented 724 (83.8%), while 129 (14.9%) were non-Saudi. Of the total participants, only 21.5% (95% CI, 18.8-24.2) had a family history of breast cancer.

Breast self examination was known to 697 (81.8%) participants whereas only 518 (61.2%) of the study subjects had knowledge of mammography. All participants had moderate knowledge of the changes that occur in breast cancer, but only 49.5% knew about the size changes, 49% were aware of heaviness under the armpit, 47% knew about touch, 45.4% had knowledge of discharge from the nipple and only 40.9% of the sample knew about the change in the shape of the nipple.

The level of knowledge of risk factors that increase susceptibility and knowledge of protective factors is shown in *Table 2*. The main sources of knowledge were television (59.4%), the internet (26.4%), magazines (55%), books (29.1%), relatives (16.3%), friends (19.2%) and the doctor (24.5%). Some of the participants had more than one source of information. Breast self examination had been performed by 41.2% (95% CI, 37.9-44.5) of the participants and 18.2% (95% CI, 15.5-20.8) had undergone mammography.

The univariate analysis showed that there was a highly statistically significant association between marital status, educational status and family history of breast cancer of the participants and knowledge of BSE, mammography and the practice of BSE and use of mammography. Married women had significantly greater knowledge and practiced BSE and mammography more than single and other women. Women with a university level education had significantly greater knowledge and practice of BSE and

Table 1: sociodemographic characteristics of study sample (n=847)

Characteristics		
Age		
Below 20	131	15.2
20-24	260	30.1
25-29	134	15.5
30-34	113	13.1
35-39	74	8.6
40-44	61	7.1
45-49	49	5.7
50+	30	3.5
Unknown	12	1.4
Marital status		
Single	433	50.3
Married	391	45.4
Divorced	26	3.0
Widow	11	1.3
Educational level		
Illiterate	10	1.2
Primary-secondary	268	31.0
University	586	67.8
Occupation*		
Students	206	29.3
Housewives	195	27.7
Health workers	71	10.1
Educational workers	115	16.3
others	117	16.6

*n=704

mammography than the women of other levels of educational status. Women with a family history of breast cancer had significantly greater knowledge and practice of BSE and mammography *Table 3* and *Table 4*.

DISCUSSION

The etiology of breast cancer is still uncertain and as a result adequate primary prevention is difficult. Hence, early detection remains the first priority for national health promotion programs. These measures include breast self-examination, which is a unique procedure in many ways because it is in-

Table 2: participants knowledge about breast cancer and its associated risk and protective factors.

	count (%)	95% Confidence intervals
Knowledge of breast cancer		
Knowledge of BSE (n=852)	697 (81.8)	79.2-84.4
Knowledge of mammography (n=847)	518 (61.2)	57.9-64.5
Heard of BSE (n=695)	569 (81.9)	79.0-84.8
Heard of mammography (n=687)	434 (63.2)	59.6-66.8
Changes that occur in relation to breast cancer (n=864):		
Size Changes	428 (49.5)	46.2-52.8
Heaviness under armpit	423 (49)	45.7-52.3
Touch	406 (47)	43.7-50.3
Discharge from nipple	392 (45.4)	42.1-48.7
Shape of nipple	353 (40.9)	37.6-44.2
Knowledge of risk factors (n=864):		
Contraceptive pills	261 (30.0)	27.1-32.9
HRT	411 (47.6)	44.3-50.9
Exposure to excess radiation	410 (47.5)	44.2-50.8
Smoking	472 (54.6)	51.3-57.9
Hereditary	338 (39.1)	35.9-42.3
Age	119 (13.8)	11.5-16.1
Nutrition	353 (40.9)	37.6-44.2
Lack of physical exercise	137 (15.9)	13.5-18.3
Irregular cycles	54 (6.3)	4.7-7.9
Race	45 (5.2)	3.7-6.7
Infertility	140 (16.2)	13.7-18.6
Knowledge of protective factors (n=864):		
Breast feeding	597 (69.1)	66-72.1
Nutrition	536 (62.0)	58.8-65.2
Exercise	322 (37.3)	34.1-40.5
Regularity of menstrual cycle	256 (29.6)	26.6-32.6
Pregnancy earlier than 40 years	217 (25.1)	22.2-28.0
Early marriage	103 (11.9)	9.8-14.0

Table 3: Association between characteristics of sample with their Knowledge and practice of breast self examination.

Characteristics of sample	Know BSE count (%)		Practice BSE count (%)	
	Yes	No	Yes	No
Marital status				
married	374 (53.8)	53 (34.2)	223 (63.7)	202 (40.5)
Single	294 (42.3)	94 (60.6)	110 (31.4)	278 (55.7)
Others	27 (3.9)	8 (5.2)	17 (4.8)	19 (3.8)
Educational level				
Primary	6 (0.9)	2 (1.4)	2 (0.6)	5 (1.0)
Intermediate	29 (4.2)	7 (4.9)	12 (3.4)	34 (6.8)
High school	153 (22.0)	53 (36.8)	64 (18.3)	145 (28.9)
University	474 (68.3)	77 (53.5)	254(72.8)	298 (59.5)
Others	32 (4.6)	5 (3.5)	17 (4.9)	19 (3.8)
Family history				
Yes	153 (22.5)	31 (20.3)	95 (27.7)	89 (18.2)
No	527 (77.5)	122 (79.7)	248 (72.3)	401 (81.8)

† Marital status, $\chi^2= 19.5$, $P<0.0001$; Educational level, $\chi^2= 25.9$, $P<0.0001$; $\chi^2= 0.245$, $P= 0.62$

• Marital status, $\chi^2= 49.3$, $P<0.0001$; Educational level, $\chi^2= 19.8$, $P<0.0001$; Family history, $\chi^2= 10.1$, $P<0.0001$

Table 4: Association between characteristics of sample with their Knowledge and practice of mammography.

Characteristics of sample	Know of mammography count (%) †		Had mammogram count (%) *	
	Yes	No	Yes	No
Marital status				
married	289 (56.0)	134 (40.8)	117 (75.9)	303 (44.7)
Single	203 (39.3)	183 (55.8)	26 (16.9)	352 (52.0)
Others	24 (4.6)	11 (3.3)	11 (7.1)	22 (3.2)
Educational level				
Primary	4 (0.8)	2 (0.6)	2 (1.3)	6 (0.9)
Intermediate	15 (2.9)	32 (9.7)	(3.9)	36 (5.3)
High school	106 (20.6)	103 (31.2)	36 (23.5)	167 (24.5)
University	362 (70.4)	185 (56.1)	100(65.3)	443 (65.1)
Others	27 (5.2)	8 (2.4)	9 (5.9)	28 (4.1)
Family history				
Yes	136 (26.7)	49 (15.3)	44 (29.5)	137 (20.5)
No	374 (73.3)	271 (84.7)	105 (70.5)	532 (79.5)

† Marital status, $\chi^2= 21.9$, $P<0.0001$; Educational level, $\chi^2= 25.9$, $P<0.0001$; Family history, $\chi^2= 13.9$, $P<0.0001$

• Marital status, $\chi^2= 62.9$, $P<0.0001$; Educational level, $\chi^2= 1.4$, $P=0.70$; Family history, $\chi^2= 5.28$, $P=0.02$

expensive, non-invasive, involves little time and physical energy, is simple and does not depend on professional help. Knowledge of breast cancer risk factors and protective factors are also very important in the health of women. Regardless of family history, women still need to be “breast aware” and to accurately identify breast symptoms in order to receive treatment as quickly as possible.

In the present study, it was found that knowledge of breast self-examination was high when compared with knowledge of mammography. In contrast, the practice of BSE was low and a high rate for having had a mammogram was observed only in women who had a family history of breast cancer. Maha and Hadi^[12] showed that 78% of 300 women had previously performed BSE, but only 42.7% had agreed to a mammography screening. This is in contrast to our finding, where it was found that 41.2% women had performed BSE and only 18.2% had mammography screening. Hadi’s study also rated the level of knowledge of methods to detect breast cancer as low and found that 3.3% of 300 of those who had a positive family history had performed BSE while 6.7% had a mammography screening. In this study, 51.6% of 184 of those who had a positive family history had performed BSE while only 24.3% had a mammography screening. The results imply that even women with a previous family history of breast cancer were not performing BSE and mammography testing.

While we expected that more women who performed BSE would have a previous family history, our results showed the opposite, which might be either the result of lack of knowledge of BSE or poor technique or it may be due to the larger sample of those participants without a previous family history. The marital status of women was useful in acquiring knowledge of BSE and its practice, which leads in knowing about mammogram also to perform mammogram.

It is generally interesting to know how education reflects on the knowledge and practice of BSE and if higher education would make women more aware of the disease. Our results revealed that women who reached a primary-secondary educational level had less knowledge and practiced BSE poorly, while women who had at least a bachelors degree had better knowledge and practice. Knowledge of breast changes that occur when disease develops was below 50% for all the symptoms.

Knowledge of risk factors is the essence of prevention. The results of our study show a low-to-moderate level of knowledge, regardless of the women’s educational status, marital status and

their family history of breast cancer. The same pattern was observed in relation to the knowledge and practice of mammography. The sources of knowledge among these women was diverse, coming from more than one resource.

In spite of the limitations encountered while conducting this study, the results suggest that knowledge of breast cancer and its associated factors was not that high, and differed in relation to the characteristics of the women. But there is a need to improve knowledge and practice of women regarding issues related to breast cancer prevention, early detection and intervention through continuous awareness programs. And these educative programs will help in preventing the disease and also help women at higher risk for developing breast cancer.

As the total sample were taken from the Ladies Kingdom Mall, a luxurious mall in the north of Riyadh city and since most of the women are from a high social class, therefore this sample cannot be generalized to the total population of Saudi Arabia. Selection bias is unavoidable in any study comparing women with high educational levels to those of lower educational levels. Highly educated women, particularly those with a family history of breast cancer, are more likely to be referred for early screening mammography and to receive a breast examination, therefore allowing their breast cancers to be diagnosed at an earlier stage than others. As a consequence, breast cancer survival in highly educated women with a previous family history should be improved due to the detection of the breast cancer at an earlier stage of the disease.

RECOMMENDATIONS

The results have shown that most women are familiar with BSE, but few perform BSE. Even those considered educated were either not aware of how to perform BSE or were simply not doing it. It is therefore of utmost importance to launch educational programs to increase public awareness of the prevalence of breast cancer, its risk factors and the importance of educating woman. Awareness should be enhanced by all available means. Important tools include news media, schools, social gatherings, hospital waiting areas, and distribution of pamphlets. The Ministry Of Health and educational facilities should play a leading role in increasing awareness. Physicians and surgeons need to make people aware of the increasing incidence of breast cancer during routine patient visits. They should use their influence to promote public education. Mobile units should provide services to people

living in remote areas. Health professionals should be required to provide mammography and genetic counseling for those who are at high risk. Pre-marital medical examination and counseling, which has been made mandatory recently in the Kingdom, is a step in the right direction. It is hoped that women

will become more aware of their health responsibility. Routine check-ups should be encouraged at the primary health care level.

Special thanks to Mr. Muneer in the Department of Family and Community Medicine for his technical assistance and data entry.

REFERENCES

1. K McPherson, C M Steel, J M Dixon. Breast cancer-epidemiology, risk factors, and genetics. *ABC of Breast Diseases* 2000; 321: 624-628.
2. Al-Fudi A, Parkin DM, Al Khogali M. Cancer in Iraq: seven-year data from the Baghdad tumor registry. In: Al Khogali M, editor. *Cancer prevention in developing countries*. London: Pregamon Press 1986: 25-34.
3. El-Harthi A. El-Harthi, Maha S. Abdel-Hadi, Thilo Doerk, Ahmed N. Badr and Joerg Schmidtk. The potential benefits of genetic testing in breast and ovarian cancer. *Saudi Med J* 1999; 20 (9): 663-670.
4. Samir S. Amr, Abdul Rahman M. Sa`di, Fazal Ilahi and Salwa S. Sheikh. The spectrum of breast diseases in Saudi Arabia Females: A 26 years pathological survey at al Dhahran Health center. *Ann Saudi Med* 1995; 15 (2): 125-132.
5. *Cancer Incidence Report*. Saudi Arabia: Ministry of Health 1994-1996.
6. *Cancer incidence report, Saudi Arabia 1997-98 (2001)*. National cancer registry, Ministry of Health, Kingdom of Saudi Arabia.
7. Dorothy Makanjoula, Khalid Murshid, Abdulkarim Elbakery, Mervet Al Saleh and Ammar Al Rikabi. Early detection of Breast Cancer: Report from King Khalid University Hospital. *Ann Saudi Med* 1996; 16(2): 139-143.
8. El-Harth A. El-Harthi, Maha S. Abdel-Hadi, Diana Steimann and Thilo Dork. BRCA1 and BRCA2 mutations in breast cancer patients from Saudi Arabia. *Saudi Med J* 2002; 23(6): 700-704.
9. Hesham A. Najjar. Breast Cancer- Is there a need for a breast unit? (Letter to the editor). *Saudi Med J* 2001; 22(5): 460-462.
10. 2001 Tumor Registry Annual Report. King Faisal Specialist Hospital and Research Center (personal communication).
11. Tawfik M. Al Tamimi, Ezzeldin M. Ibrahim. Abdel Wahab M. Ibrahim, Adnan A. Al Bar, Suliman A. As-suhaimi, Gabriel S. Gabriel, Adel M. Mishriky, Hassan Y. Al Idrissi, Mohammad O. Al Sohaibani and Mohammed H. Al Sibai. Cancer In the Eastern region of Saudi Arabia: A population-Based study (1987-1988). *Ann Saudi Med* 1997; 17(1): 53-65.
12. Maha S.A. Abdel Hadi. Breast Cancer Awareness among Health Professionals.

THE KNOWLEDGE OF BREAST CANCER AMONG YOUNG SAUDI FEMALES*

Wafa A. Sait, MBBCh, Arab Board, Samia M. Al-Amoudi, Arab Board, Dalal A. Tawtai, MBBCh, Hassan S. Abduljabbar, MD, FRCS.

ABSTRACT

الأهداف: تقييم مستوى المعلومات التي تمتلكها طالبات المدارس السعوديات حول سرطان الثدي.

الطريقة: أُجريت هذه الدراسة الاستفتائية في مستشفى جامعة الملك عبدالعزيز، جدة، المملكة العربية السعودية وذلك خلال الفترة من أبريل إلى يونيو 2009م. شارك في هذه الدراسة 500 طالبة من مختلف المدارس الثانوية بجدة. لقد قمنا بتحديد مواقع المدارس في مختلف أنحاء مدينة جدة باستخدام الخريطة. ومن ثم قمنا بتوزيع الاستفتاء الذي يحتوي على مجموعة من الأسئلة التفصيلية حول سرطان الثدي وذلك بعد أخذ الترخيص من وزارة التربية والتعليم. وقد تدرّب مجموعة من المتطوعين على كيفية توزيع أوراق الاستبيان وجمعها. وبعد ذلك تم تحليل نتائج الاستبيان إحصائياً. لقد تم استخدام نتائج الاستفتاء من أجل بناء قاعدة للبرنامج التثقيفي حول هذا المرض.

النتائج: أظهرت نتائج التحليل لما يعادل 337 من أوراق الاستبيان بأن مستوى المعلومات التي تمتلكها طالبات المدارس والجامعات محدودة وغير كافية، إلا أنه تم ملاحظة اهتمام الطالبات وحماسهم نحو معرفة المزيد عن سرطان الثدي وكيفية الوقاية منه.

خاتمة: أشارت الدراسة إلى أن تدني مستوى المعلومات التي يمتلكها الجيل الجديد قد يكون عائقاً أمام برامج الفحص والتشخيص المبكر لسرطان الثدي. ولهذا فنحن بحاجة إلى زيادة الوعي حول هذا المرض وذلك من خلال المحاضرات، والحلقات الدراسية، وورشات العمل، وبرامج التدريب.

* Saudi Med J. 2010, Vol. 31(11)

Objectives: To assess the breast cancer knowledge level of Saudi female school students.

Methods: A detailed questionnaire on cancer breast was designed with all the needed information. Using a map of the Jeddah area of Saudi Arabia, schools were identified in each area and permission was sought from the Ministry of Education to distribute the questionnaire to the students. A team of volunteers was instructed on how to distribute and collect the questionnaires. The collected questionnaires were then statistically analyzed. This Pilot study of 500 students was performed in King Abdul-Aziz University Hospital using high schools from the city of Jeddah between April and June 2009. The results were used to build up a base for designing a community educational program.

Results: Analysis of the data from 337 questionnaires from high school and college students showed that the level of knowledge of young females on breast cancer is limited. However, it also indicated that the students are very enthusiastic to learn about cancer breast, and its prevention.

Conclusion: The limited knowledge level of breast cancer in the younger generation might be an obstacle to screening programs and early diagnosis. Awareness programs should be developed including lectures, seminars workshops, and on hands training.

Breast cancer remains the most commonly diagnosed cancer in women, and it is the second leading cause of cancer related death.^[1] Worldwide, more than a million women are diagnosed with breast cancer every year.^[2] Breast cancer incidence rates vary considerably, and have shown a marked geographical variation, with the highest rates in western countries, and lowest in Asian and African countries.^[3] The breast cancer incidence is increasing in most regions; the changes are greatest in areas where rates were previously low.^[4] Environmental factors might be the cause for the global variations. It was found that women from Japan that emigrate to the USA ultimately lose the advantage of low risk, and within one to two generations they develop the same risk profile as USA residents.^[4] For primary prevention of breast cancer, women need to be adequately informed of risk factors and risk reduction strategies for breast cancer.^[5] Low cancer awareness contributes to delay in presentation for cancer symptoms, and may lead to a delay in cancer diagnosis.^[6] Breast selfexamination detects new breast

cancers in high-risk women undergoing screening mammogram, clinical breast examination, and yearly breast MRI.^[7] The aim of this research is to assess the breast cancer knowledge level of Saudi female school students, which will help in designing breast cancer awareness education programs for the younger generations.

METHODS

This pilot study was carried out at King Abdulaziz University Hospital, Jeddah, Saudi Arabia between April and June 2009, and the research proposal was approved by the Research Ethical Committee of the institute. A map of Jeddah was used to locate the schools in each area, and permission was sought from the Ministry of Education to distribute the questionnaires to the students. A team of 4 volunteers' and 2 physicians were instructed on how to distribute and collect the questionnaire. Verbal informed consent was taken from all the students prior to participating and completing the questionnaire.

The detailed questionnaire on breast cancer included 9 questions in the first section: name, age, age of menarche, level of education, social status, family history, what is a mammogram, if there are any symptoms of breast disease, and if there any surgical procedures in the breast. The second part of the questionnaire comprised 6 questions on the risk factors of breast cancer, including knowledge of oral contraceptive pills, radiation, smoking, fatty foods, family history of ovarian and colonic cancer. This was followed by 3 questions on self-breast examination, like how and when to carry out a self-examination, and if they would like to receive training on how to carry this out. Finally, there were 5 general questions including relationship of breast-feeding, breast brassiere, if there is any treatment for breast cancer, does it cause death, and should the diagnosis be secret.

The questionnaires were collected and the data were entered and analyzed using the Statistical Package for Social Sciences version 15 (SPSS Inc, Chicago, IL, USA). Percentages of the different variables were calculated, risk factors and knowledge about cancer breast was analyzed, and the results were used to build up a base for designing a community educational program.

RESULTS

Out of 500 students, only 337 answered. On analysis of 337 questionnaires, 21% were college students, and 79% were high school level, 99% were single and only 1% were married. Age ranged from

12-18 years with a mean of 16.2 ± 1.5 , and the age of menarche between 9-16 years with a mean of 12.6 ± 1.2 years. Of the 337 students, 9.8% admitted that they have a family history of breast cancer, 83.9% answered that they have no family history of breast cancer, and 8.3% did not know. On direct questioning on mammograms, only 30% knew what this was, and 70% did not know. Only 18.4% had complaints regarding their breasts, such as mastalgia, pain, lumps, or their breasts were getting bigger, and only 1.8% reported previous surgery, mainly aspiration, or biopsy. *Table 1* summarizes the results of exploring the level of knowledge of certain risk factors, namely, oral contraceptive pills (OCP), radiation, smoking, and fatty food. On direct question regarding carrying out self-breast examination, 61.1% answered yes and 39.9% answered no. Surprisingly approximately 55% answer that it has to be carried out after the monthly period and not before as recommended, and 72.1% of the students were very enthusiastic about receiving a training course to learn how to perform self-breast examination. Only 2% of students reported a family history of ovarian or colonic cancer. *Table 2* summarizes the results of questions regarding knowledge of breast cancer prevention and treatment.

DISCUSSION

It seems that the younger Saudi generation has limited information and knowledge of cancer breast, and it is well known that low cancer awareness contributes to delays in presentation of cancer symptoms, and subsequent diagnosis.^[6] Approximately 40% of the study population reported a lack of knowledge regarding breast self examination. Although, breast self examination has not been proven to be effective in early detection or in lowering mortality from breast cancer,^[7] it is a good way to raise awareness regarding the appropriate age to start screening, namely, periodic clinical breast examination and mammographic screening. Screening mammography is the only modality proven by randomized clinical trials to allow early detection resulting in overall lower mortality.^[8] In our study, only 30% of our students knew of the mammogram. It is also known that mammographic screening is effective not only in women aged 50 years or more, but also in those aged less than 50 years.^[8] The risk factors for breast cancer are well known, such as female gender, older age, and the older the age the greater the risk of breast cancer, a positive family history of breast cancer, being exposed to large amounts of radiation, such as very frequent spinal x-rays for scoliosis or treatment for Hodgkin's dis-

Table 1: Results of the questions related to knowledge of breast cancer risk factors.

Risk factor	Yes	No	Don't Know	Missing**
OCP	32.6	26.7	39.2	1.5
Radiation	60.8	13.6	24.0	1.5
Smoking	65.0	15.1	19.3	0.3
Fatty food	48.4	14.2	35.6	1.8

**missing means did not answer, OCP - oral contraceptive pill

Table 2: Results of the questions related to knowledge of breast cancer prevention and treatment.

Variable	Yes	No	Don't Know	Missing**
Breast feeding	68.8	6.2	22.3	2.7
Breast brassiere	44.2	19.9	31.5	4.5
Any treatment	77.7	2.4	16.0	3.9
Can cause death	0.6	32.6	63.2	3.6
Should you talk about breast cancer	3.0	85.5	5.6	5.9

**missing means did not answer.

ease at a young age, a personal history of breast or ovarian cancer, being overweight after menopause, or gaining weight as an adult and current or recent use of birth control pills.^[9-17] A recent meta-analysis of 54 studies relating OCP use to breast cancer found that women who are currently using combined OCP or have used them in the past 10 years are at a slightly increased risk (relative risk 1.07-1.24) compared with never users.^[14]

A similar study carried out by Millat in 2000,^[18] concluded that female secondary-school students in Jeddah had very little knowledge of the presentation of breast cancer and its risk factors. Students were also not familiar with breast self-examination. The study indicated the necessity for a health education program on risk factors, early signs, and methods of diagnosis of breast cancer for this group of easily targeted young women.^[18]

In conclusion, our results indicate the limited information about cancer breast in the younger generation, which might be an obstacle to screening programs and early diagnosis. However, we also

found that the younger generation was very enthusiastic to learn more about breast cancer. An awareness program has to be developed including lectures, seminars, workshops, and hands on training. To improve the medical knowledge of young women seeking education, this information could be introduced in the basic curriculum in high school.

ACKNOWLEDGMENT

This study was sponsored by the Scientific Chair of Sheikh Mohammad H. Al-Amoudi for Breast Cancer. We would like to thank Mr. Abdulkareem Al-Haqeel, Mrs. Nadiah Thabet, Mrs. Noor S. Ba Qader, Dr. Eman M. Kasim, and Mrs. Somaya Abdulwahab for their help in obtaining the necessary permission to perform this study, and their help in distributing, and collecting the questionnaires.

REFERENCES

1. American Cancer Society. Breast Cancer Facts & Figures 2009-2010. Atlanta: American Cancer Society; 2009. [Updated 2010 July 25; Accessed 2009 November 3] Available from: <http://www.cancer.org/research/cancerfactsfigures/breastcancerfactsfigures/breast-cancer-facts--figures-2009-2010>.
2. Ferlay J, Bray F, Pisani P, Parkin DM, editors. Globocan 2002: Cancer Incidence, Mortality and Prevalence Worldwide, Version 2.0. Second edition: Lyon (France): IARC Press; 2004.
3. Jemal A, Siegel R, Ward E, Hao Y, Xu J, Thun MJ. Cancer statistics, 2009. *CA Cancer J Clin* 2009; 59: 225-249.
4. Anderson BO, Shyyan R, Eniu A, Smith RA, Yip CH, Bese NS, et al. Breast cancer in limited-resource countries: an overview of the Breast Health Global Initiative 2005 guidelines. *Breast J* 2006; 12 Suppl 1: S3-S15.
5. Liebens F, Aimont M, Beauraing F, Carly B, Cusumano P, Rozenberg S. What information do public organizations provide to Belgian women on primary prevention of breast cancer? *Eur J Cancer Prev* 2010; 19: 68-70.
6. Austoker J, Bankhead C, Forbes LJ, Atkins L, Martin F, Robb K, et al. Interventions to promote cancer awareness and early presentation: systematic review. *Br J Cancer* 2009; 101: S31-S39.
7. Wilke LG, Broadwater G, Rabiner S, Owens E, Yoon S, Ghate S, et al. Breast self-examination: defining a cohort still in need. *Am J Surg* 2009; 198: 575-579.
8. Gøtzsche PC, Nielsen M. Screening for breast cancer with mammography. *Cochrane Database Syst Rev* 2006; 4: CD001877.
9. Morimoto T, Sasa M, Yamaguchi T, Kondo H, Akaiwa H, Sagara Y. Breast cancer screening by mammography in women aged under 50 years in Japan. *Anticancer Res* 2000; 20: 3689-3694.
10. Tretli S. Height and weight in relation to breast cancer morbidity and mortality. A prospective study of 570,000 women in Norway. *Int J Cancer* 1989; 44: 23-30.
11. Reeves GK, Pirie K, Beral V, Green J, Spencer E, Bull D, et al. Cancer incidence and mortality in relation to body mass index in the Million Women Study: cohort study. *BMJ* 2007; 335: 1134.
12. Jee SH, Yun JE, Park EJ, Cho ER, Park IS, Sull JW, et al. Body mass index and cancer risk in Korean men and women. *Int J Cancer* 2008; 123: 1892-1896.
13. Friedenson B. Oral contraceptives and the risk of breast cancer. *N Engl J Med* 2002; 347: 1448-1449.
14. Fentiman IS. Oral contraceptives, hormone replacement therapy and breast cancer. *Int J Clin Pract* 2002; 56: 755-759.
15. Ahn J, Schatzkin A, Lacey JV Jr, Albanes D, Ballard-Barbash R, Adams KF, et al. Adiposity, adult weight change, and postmenopausal breast cancer risk. *Arch Intern Med* 2007; 167: 2091-2102.
16. Sweeney C, Blair CK, Anderson KE, Lazovich D, Folsom AR. Risk factors for breast cancer in elderly women. *Am J Epidemiol* 2004; 160: 868-75.
17. Krebs EE, Taylor BC, Cauley JA, Stone KL, Bowman PJ, Ensrud KE. Measures of adiposity and risk of breast cancer in older postmenopausal women. *J Am Geriatr Soc* 2006; 54: 63-69.
18. Milaat WA. Knowledge of secondary-school female students on breast cancer and breast self-examination in Jeddah, Saudi Arabia. *East Mediterr Health J* 2000; 6: 338-343.