

ORIGINAL ARTICLES

Changing trends of histopathology in childhood nephrotic syndrome in western Saudi Arabia

Jameela A. Kari

ABSTRACT

Objectives: It is widely accepted that minimal change nephrotic syndrome is the most common cause of nephrosis in children. Recent studies have demonstrated an increasing incidence of focal segmental glomerulosclerosis in adults and children. This study was conducted to analyze the trend of histopathologic subtypes in idiopathic nephrotic syndrome in the western area of the Kingdom of Saudi Arabia.


Methods: To determine the possible changes in the etiology of childhood nephrosis, the clinical charts of 46 pediatric patients diagnosed with idiopathic nephrotic syndrome between 1997 and 2001, group A, were reviewed. Results were compared with our initial published data of 132 patients with idiopathic nephrotic syndrome diagnosed between 1983 and 1992, group B.

Results: There was no difference in the percentage of biopsies between group A (43%) and group B (31.8%), $P=0.9$. There was a decline in the incidence of minimal change nephrotic syndrome in recent years. Even if we assume that patients without a histological diagnosis had minimal change nephrotic syndrome, presumptive minimal change nephrotic syndrome, the total incidence of minimal change nephrotic syndrome (biopsy proven + presumptive) in group A, was only 65% compared to 79.5% in group B, $P=0.02$. The incidence of focal segmental glomerulosclerosis was significantly greater in biopsies performed in the recent period, group A, 35% versus group B, 16.7%, $P=0.05$, and in the total number when we included the presumptive minimal change nephrotic syndrome, group A, 15.2% and group B, 5.3%, $P<0.0001$. Similarly we found an increased incidence of membranoproliferative glomerulonephritis in total patients when we included the presumptive minimal change nephrotic syndrome, group A, 13.0% and group B, 5.3%, $P=0.02$.

Conclusions: There is a shift toward an increasing prevalence of focal segmental glomerulosclerosis and membranoproliferative glomerulonephritis over the years in the western area of the Kingdom of Saudi Arabia. These findings may have significant implications in the management of childhood nephrotic syndrome

Saudi Medical Journal 2002; Vol. 23 (3):

Nephrotic syndrome (NS) is a clinical manifestation of different histopathologic sub-types. Based on histopathological findings on renal biopsy, primary NS is further classified into minimal change nephrotic syndrome (MCNS), focal segmental glomerulosclerosis (FSGS), mesangioproliferative glomerulonephritis (MePGN), membranoproliferative glomerulonephritis (MPGN), membranous



glomerulonephritis (MGN) and focal and global glomerulosclerosis.¹ The International Study of Kidney Disease in Children (ISKDC) reported in the 1970s that MCNS was the most common histological lesion in renal biopsy from children with idiopathic nephrotic syndrome (INS).^{2,3} A similar study carried out by White et al⁴ in the United Kingdom at the same time reported an incidence of 76.5% and 8.3% of MCNS and FSGS. We have reported the same observation in our previous study of histopathological lesions in children with NS presented between 1983 and 1992.⁵ We have reported MCNS in 100% of nephrotic cases and showed like others^{2-4,6} that FSGS was a rare cause of nephrosis in children, observed in only 5.3% of the total patients and in 10% of the biopsies. More recent studies in adults and children showed changing trends of histopathology in adulthood^{7,8} and childhood NS⁹⁻¹¹ with increasing incidence of FSGS during the last 20 years, accompanied by significant decrease in the incidence of MCNS. We report the changing trend in the histopathological lesions in INS syndrome in the Western area of Kingdom of Saudi Arabia. We have retrospectively reviewed all the children presented with INS to King AbdulAziz University Hospital (KAUH) between 1997 and 2001 and compared the results with our previously published data. King AbdulAziz University Hospital is located in a racially mixed population of a major urban area in the Western area of KSA. The demographic data and the histopathological findings were reviewed.

Methods. A retrospective study was performed from January 1997 to June 2001 at KAUH, involving 46 patients aged up to 14 years with childhood NS. The results were compared with our published data of 132 nephrotics presented at the same institution between 1983 and 1992.⁵ These patients comprised a racially mixed population of a major urban city in KSA located near to the city of Makkah where expatriates are common. Most had been treated elsewhere with other physicians and referred to our unit for management. Patients with congenital nephrotic syndrome (CNS) or with systemic illness such as systemic lupus erythematosus, Henoch-Schönlein purpura, sickle cell anemia, malignancies, metabolic disorders or hepatitis were excluded. Indications for biopsy included steroid resistance, an age older than 8 years, unusual presentation such as significant elevation of serum creatinine or gross hematuria and prior cyclosporin therapy in frequent relapsers and steroid dependent nephrotics. The biopsy specimens were evaluated histopathologically by light, immunofluorescent and electron microscopy (EM). Histopathological findings were interpreted by the same pathologists in accordance with similar criteria as applied in our previous study. Adequacy of biopsy was defined as the presence of at least 5 glomeruli in the specimen on light microscopy.¹² Minimal change disease (MCD) was characterized by the absence of any conspicuous abnormality on light microscopy. Mesangioproliferative glomerulonephritis was labeled in the presence of diffuse mesangial hypercellularity lacking immune deposits. Focal segmental glomerulosclerosis was characterized by the presence of at least one glomerulus showing a segmental area of sclerosis with or without accompanying tubular atrophy and interstitial fibrosis. Membranoproliferative glomerulonephritis was labeled in the presence of immune cellular proliferation on light microscopy and immune deposition on EM. In the absence of atypical presentation of childhood NS, our practice is to treat such patients with corticosteroids. The group of patients who readily respond to prednisolone (56.5%) is presumed to have MCNS. Renal biopsy in such patients was not performed. All patients were treated initially with corticosteroids, prednisolone, at a dose of 60 mg/m²/day (maximum 80 mg/day) for 4 w

Nine patients received levamisole and 11 patients were treated with cyclophosphamide. Cyclosporin was used in 4 patients with FSGS, 2 steroid dependent and 2 steroid resistant.

Definitions. Nephrotic syndrome was defined as the association of heavy proteinuria and hypoalbuminemia. The response to therapy was classified according to the definition from the British Pediatric Nephrology Association: a. steroid sensitive - complete resolution of proteinuria within 4 weeks of prednisolone therapy, b. steroid resistant - failure to respond after 4 weeks prednisolone 60 mg/m²/day, c. steroid dependence - recurrence of nephrosis when the dose of corticosteroids is reduced or within 2 weeks after discontinuation of therapy, d. frequent relapses - 2 or more episodes of nephrosis within 6 months of the initial response or 4 or more within any month period. Congenital NS was defined as NS presented at less than 3 months of age.

Statistical analysis. Data from groups of patients are reported as mean ± standard deviation (SD) or as percentage (frequency of observation in a particular group). The statistical analysis was performed by Z test and t-test assuming equal variance was used to compare between the ages of different groups. Statistical significance was defined as a P < 0.05.

Results. Our populations were presented by one major ethnic group; Arab [N=37 (80%), 20 of them were Saudis (43%)] and 3 minor ethnic groups: Asians (Indians and Pakistanis, N=5); Africans (N=2) and 2 patients were from the Far East (Indonesian and Filipino). The mean ± SD age of our population (group A) at first presentation of NS was 3.8 ± 2.8 (range 0.7 to 14 years) compared to 6.2 ± 3.7 in group B. Three patients with CNS were excluded from the study in group A and 10 patients in group B. All patients of the study were seronegative for hepatitis surface antigen (HbsAg) and antinuclear antibodies (ANA). The fraction of patients who underwent biopsy in the 2 groups was similar, group A 20 of 46 (43%) versus group B, 42 of 132 (31.8%), P=0.05. In group A, children with INS who underwent biopsies, FSGS was the most common histopathological subtype, occurring in 7 children (35%), followed by MPGN in 6 children (30%). Other subtypes included MesPGN in 4 patients (20%) and MCD in 3 patients (15%), Table 1. If we assumed that all children that were not biopsied had MCD (presumptive MCD), the total incidence of MCNS (presumptive + biopsy proven) was 65%, which is lower than that of group B (79.5%), Z= 2.02, P=0.02, Table 2. The prevalence of FSGS was significantly higher in-group A (35% of the biopsies versus 16.7% in group B, P=0.05), Table 3. The increase in fraction of patients with FSGS was significant in group A, even if all patients including those who did not undergo biopsies were considered, group A, 7 of 46 (15.2%) versus group B, 7 of 132 (5.3%), P=<0.0001, Table 2. There was a non-significant increase in the MPGN in biopsied patients, group A 6 of 20 (30%) versus 7 of 42 (16.7%), P=1.2, Table 3. However, when we included patients who did not have renal biopsies and a presumptive diagnosis of MCD, the increase in MPGN was significant, group A 6 of 46 (13%) versus group B 7 of 132 (5.3%), P=0.02, Table 2. We thought about the effect of the racial origin, however, when we studied patients from Arab origin only 37 patients, after excluding Asian and African patients, we found similar results. Fifteen patients (40.5%) had a renal biopsy. Of those 6 patients (40%) had FSGS, 6 (40%) had MPGN and 3 (20%) had MesPGN. The Saudi

patients, 20 children, as a sub-group had the same distribution of the whole group or the Arab group. Seven had a renal biopsy, 3 (42.8%) had FSGS, (42.8%) had MPGN and 1 (14.2%) had MesPGN, Table 4. We do not have racial fractions of our previous study but they were from the same population. Familial steroid resistant nephrotic syndrome (SRNS) was observed in 3 Arab siblings from Yemeni origin with MPGN. Another patient with SRNS due to MPGN had a family history of a sibling who died from chronic renal failure complicated SRNS. The mean \pm SD of total (presumptive and biopsy proven) MCD was 3.6 ± 2.2 , range 0.9-11, years. There was no difference between presenting age (mean \pm SD) of MPGN patients; 2.8 ± 3.1 , range 0.8-5 years and MCNS patients, $P=0.6$. However, FSGS patients had an older age at presentation 6.14 ± 4.7 , range 2-14, years, $P=0.03$. Ten patients were steroid resistant, 8 never responded to steroids and 2 were initially steroid dependent frequent relapses and changed to resistant after 6 months and 3 years from time of first presentation. Twenty-two were frequent relapsers (61% of steroid sensitive patients) and 14 patients were infrequent relapsers. There was no difference in the percentage of initial steroid resistance between group A and group B (11%), Z value=1 and $P=0.15$. Cyclophosphamide was used in 10 children, 3 steroid resistant and 8 steroid dependent. All patients with SRNS had a renal biopsy before the cyclophosphamide and 6 of the steroid dependent patients (75%). All except one were found to have a non-minimal change in histopathology.

Discussion. We have found an increasing incidence of FSGS in Arab children who presented with INS over the last decade. Our results showed that FSGS was the most common histopathologic sub-type in our patients who underwent renal biopsy in the recent years, compared to MCD in those who were biopsied between 1983 and 1992. This observation is similar to the previous reports from different parts of the world. Bonilla-Felix et al reported an increased incidence of FSGS in American children (23% before 1990 versus 47% after 1990), particularly in those from the African origin in whom FSGS was found to be the most common cause of NS.⁹ Similarly, Srivastava et al found a high incidence of FSGS in American children with reciprocal decline in the incidence of MCD in recent years.¹⁴ Gulati et al reported an increased incidence of FSGS in Northern and Eastern Indian populations from 20% between 1990 and 1992 to 47% between 1992 and 1996.¹⁰ Similar to the authors in the above studies we were careful to exclude any known secondary etiology of FSGS and believe that there is a true universal increase in the incidence. The increased incidence of FSGS in our study is in agreement with that reported in adult and pediatric populations from other countries.⁷⁻¹⁴ There is no similar reported study in adult or pediatric populations from the Arab world.

In the 1980s AbdulRahman et al reported MesPGN as a frequent cause of minimal change NS in children from the central area of KSA and they reported FSGS as a rare cause.¹⁵⁻¹⁶ In 1990 Mattoo et al reported the same observation on the same population.¹⁷ However, they reported a high prevalence of FSGS (39%) in biopsies in children with NS. Al-Raheed et al reported FSGS to be the cause of childhood INS in 24% of the biopsies performed between 1982 and 1994 but they did not study the increased incidence of FSGS over the years.¹⁸ Mitwalli et al found that FSGS was the most common glomerular disease in 186 biopsies in adult patients between 1989 and 1994.¹⁹ Ethnicity seems to play a critical role in the epidemiology

NS. Minimal change nephrotic syndrome which is usually steroid responsive nephrotic syndrome (SSNS) is more common in Asian children with an incidence of 16 per 100,000 children,²⁰ compared to 2-7 per 100,000 European children per year.²¹ While FSGS is more common in black children and the most common cause of NS in Africans and Americans,⁹ the reported incidence in Arab children is variable between 6 and 11.6 per 100,000 population.² We have considered the effect of the mixed racial composition of our population in the increasing incidence of FSGS, however we have found a similar increasing incidence of FSGS in Arab patients and in Saudi patients when were studied separately. The observed trend is unlikely to be caused by a change in the referral patterns, as there was no difference in the proportion of steroid resistance children in our patient populations over the 2 periods. Furthermore, the mean age of presentation over the recent years was similar to those in group B, while FSGS patients had an older age at presentation. Our findings support the notion that there is a global trend of increasing FSGS incidence in childhood NS.⁹⁻¹¹ Unfortunately, as the etiology of FSGS is unknown, it is difficult to postulate a potential mechanism for this increasing incidence. Previous studies suggested that it could be secondary to environmental factors including infectious factors related to viruses, chemical exposure or many factors together.⁹⁻¹¹ We have found also an increasing incidence of MPGN in recent years. This observation is different from previous studies.^{9-11,14} Membranoproliferative glomerulonephritis is a rare cause of childhood INS,^{24,3-4} we have found that 16.7% of the performed biopsies in group B and 30% in group A were MPGN.⁵ This is in contrast to the paucity of MPGN as a cause of INS in other Saudis¹⁶⁻¹⁸ and in international studies.¹⁶⁻¹⁸ Furthermore the increasing incidence has not been reported before. Membranoproliferative glomerulonephritis was reported as the predominant histological lesion seen in childhood nephrotic syndrome in Ibadan, Nigeria where MCN remains a rare cause of NS.²⁵ Membranoproliferative glomerulonephritis was found to be more prevalent (23%) than FSGS (19%) in the children with nephrotic syndrome in Auckland, New Zealand.²⁶ The observed increase in MPGN in recent years could be explained by environmental factors such as antigen-driven mechanism: infective antigen as well as food and other allergens. Alternatively, genetic background could play a role, as all MPGN patients in our study were Arabs in origin and familial MPGN was noted in 2 families. Membranoproliferative glomerulonephritis was most likely the cause of NS in the dead sibling as histopathological findings in siblings with familial NS show close to a 100% concordance rate.²⁷ Familial MPGN has been reported previously and genetic cause is the most likely predisposing factor in those families.²⁸⁻²⁹ We have also observed that the age of presentation to renal biopsy for MPGN was younger in our population compared to previous studies.^{30,31}

In conclusion, there is an increasing incidence of FSGS and to a lesser extent the MPGN in children presenting with INS in the western province of KSA. The changing trend of histopathologic patterns has profound prognostic significance. Our finding is in agreement with the recommendation of performing renal biopsies on children with INS who are steroid dependent,^{9,11} in addition to those who are steroid resistant particularly before starting cytotoxic medications. This approach will be beneficial in the management of those patients and it will help to solve the mysteries of the disease processes involved.

From the Department of Pediatrics, King AbdulAziz University Hospital, Jeddah, Kingdom of Saudi Arabia.

Received 21th July 2001. Accepted for publication in final form 28th October 2001.

Address correspondence and reprint request to: Dr. Jameela Kari, Associate Professor, Department of Pediatrics, King AbdulAziz University Hospital, P.O. Box 80215, Jeddah 21589, Kingdom of Saudi Arabia. Tel. +996 (2) 55677904 ext. +996 (2) 6743781. E-mail: jkari@doctors.org.uk

References

1. Nash MA, Edelman CM Jr, Bernstein J, Barnett HL. The nephrotic syndrome. In: Edelman CM Jr, editor. Pediatric Kidney Disease. 2nd ed. Boston: Little Brown; 1992. p. 1247-1266.
2. Nephrotic syndrome in children: prediction of histopathology from clinical and laboratory characteristics at time of diagnosis. A report of the International Study of Kidney Disease in Children. *Kidney Int* 1978; 13: 159-165.
3. Churg J, Habib R, White RH. Pathology of the nephrotic syndrome in children: a report for the International Study of Kidney Disease in Children. *Lancet* 1970; 760: 1299-1302.
4. White RH, Glasgow EF, Mills RJ. Clinicopathological study of nephrotic syndrome in childhood. *Lancet* 1970; 1: 1353-1359.
5. El Sheikh S, Andigani A, Saggaf H, Kari J, Al Shohaib S, Nassef O et al. Clinicopathological study of primary nephrotic syndrome in the western area of Saudi Arabia. *Journal of King Abdulaziz University Medical Science* 1994; 34: 33-34.
6. Hogg RJ, Silva FG, Berry PL, Wenz JE. Focal segmental glomerulosclerosis in children with idiopathic nephrotic syndrome. A report of the Southwest Pediatric Nephrology Study Group. *Kidney Int* 1985; 27: 442-449.
7. Haas M, Spargo BH, Coventry S. Increasing incidence of focal-segmental glomerulosclerosis among adult nephropathies: a 20-year renal biopsy study. *Am J Kidney Dis* 1995; 26: 740-750.
8. DAgati V. The many masks of focal segmental glomerulosclerosis. *Kidney Int* 1994; 46: 1223-1241.
9. Bonilla-Felix M, Parra C, Dajani T, Ferris M, Swinford RD, Portman RJ et al. Changing patterns in the histopathology of idiopathic nephrotic syndrome in children. *Kidney Int* 1999; 55: 1885-1890.
10. Gulati S, Sharma AP, Sharma RK, Gupta A. Changing trends of histopathology in childhood nephrotic syndrome. *Am J Kidney Dis* 1999; 33: 1000-1005.

646-650.

11. Conley SB. More questions than answers. *Kidney Int* 1999; 55: 2072-
12. Oberholzer M, Torhorst J, Perret E, Mihatsch MJ. Minimum sample size for kidney biopsies for semiquantitative and quantitative evaluation. *Nephron* 1983; 34: 192-195.
13. British Association for Paediatric Nephrology and Research Unit, Royal College of Physicians. Consensus statement on management and audit protocol for steroid responsive nephrotic syndrome. *Arch Dis Child* 1994; 70: 151-
14. Srivastava T, Simon SD, Alon US. High incidence of focal segmental glomerulosclerosis in nephrotic syndrome of childhood. *Pediatr Nephrol* 1993; 13: 13-18.
15. Abdurrahman MB, Elidrissy AT. Childhood renal disorders in Saudi Arabia. *Pediatr Nephrol* 1988; 2: 368-372.
16. Abdurrahman MB, Elidrissy AT, Shipkey FH, Al Rasheed S, Al Mugeireh S. Clinicopathological features of childhood nephrotic syndrome in Saudi Arabia. *Ann Trop Paediatr* 1990; 10: 125-132.
17. Mattoo TK, Mahmood MA, Al Harbi MS. Nephrotic syndrome in Saudi children: clinicopathological study of 150 cases. *Pediatr Nephrol* 1990; 4: 519.
18. Al Rasheed SA, Al Mugeiren MM, Al Salloum AA, Al Sohaibani MO. Childhood renal diseases in Saudi Arabia. A clinicopathological study of 167 cases. *Int J Nephrol* 1996; 28: 607-613.
19. Mitwalli AH, Al Wakeel JS, Al Mohaya SS, Malik HG, Abu-Aisha H, Hassan OS et al. Pattern of glomerular disease in Saudi Arabia. *Am J Kidney Dis* 1997; 27: 797-802.
20. Sharples PM, Poulton J, White RH. Steroid responsive nephrotic syndrome more common in Asians. *Arch Dis Child* 1985; 60: 1014-1017.
21. Chesney RW. The idiopathic nephrotic syndrome. *Curr Opin Pediatr* 1999; 11: 158-161.
22. Zaki M, Helin I, Manandhar DS, Hunt MC, Khalil AF. Primary nephrotic syndrome in Arab children in Kuwait. *Pediatr Nephrol* 1989; 3: 218-220.
23. Elzouki AY, Amin F, Jaiswal OP. Primary nephrotic syndrome in Arab children. *Arch Dis Child* 1984; 59: 253-255.
24. Davis AE, Schneeberger EE, Grupe WE, McCluskey RT. Membranoproliferative glomerulonephritis (MPGN type I) and dense deposit disease (DDD) in children. *Clin Nephrol* 1978; 9: 184-193.

25. Asinobi AO, Gbadegesin RA, Adeyemo AA, Akang EE, Arowolo FA, Abiodun SO, et al. The predominance of membranoproliferative glomerulonephritis in childhood nephrotic syndrome in Ibadan, Nigeria. *West Afr J Med* 1999; 18: 203-206.
26. Simpson AK, Wong W, Morris MC. Paediatric nephrotic syndrome in Auckland, New Zealand. *J Paediatr Child Health* 1998; 34: 360-362.
27. Moncrieff MW, White RH, Glasgow EF, Winterborn MH, Cameron JS, Oggel CS. The familial nephrotic syndrome II. A clinicopathological study. *Clin Nephrol* 1973; 1: 220-229.
28. Bogdanovic RM, Dimitrijevic JZ, Nikolic VN, Ognjanovic MV, Rodic BD, Slavkovic BV. Membranoproliferative glomerulonephritis in two siblings: report and literature review. *Pediatr Nephrol* 2000; 14: 400-405.
29. Bogdanovic RM, Dimitrijevic JZ, Nikolic VN, Ognjanovic MV, Rodic BD, Slavkovic BV. [Familial membranoproliferative glomerulonephritis] Familijna membranoproliferacioni glomerulonefritis. *Srp Arh Celok Lek* 1999; 127: 166-167.
30. Yalcinkaya F, Ince E, Tumer N, Ekim M. The correlation between the clinical, laboratory and histopathological features of childhood membranoproliferative glomerulonephritis and response to treatment. *Turk J Pediatr* 1992; 34: 137-144.
31. Kari JA, Bamashmous H, Lingawi S, Al Sabban E, Akhtar M. Infantile Nephrotic Syndrome and Congenital Glaucoma. *Pediatr Nephrol* 2001; 16: 890-897.
-